

**Theodore B. Shybut, MD FAANA FAAOS**  
Southern California Orthopedic Institute  
Knee, Shoulder, Elbow Arthroscopy &  
Reconstructive Surgery  
Burbank, Valencia / Santa Clarita, Van Nuys  
@shybutMD / [www.shybutmd.com](http://www.shybutmd.com)



## **ACL Recon + Protected Meniscal Repair Protocol (Meniscal Root / Bucket Handle / Large or Unstable Tears)**

### Tips for Successful Recovery

- 1. Surgical pre-cautions:** Do not change bandages unless instructed by physician. Wear compression hoses on operative limb until crutches are discharged. If you suspect a DVT, contact Dr. Shybut's office immediately at 661-290-5473 or refer to ED immediately. If patient has reactive effusion that does not improve with rest, ice, and compression, contact Dr. Shybut's office.
- 2.** Begin stretching extension ROM on day one. Achieve full extension ROM by week 2. If not achieved by end of week 4, notify the physician's office.
- 3.** Address quad activation early and focus on isolation of quadriceps activation. Use surface emg, NMES, and cueing to isolate quadriceps. Be aware of co-contracting from hamstrings, and ensure proper form. Do not progress to standing activities if patient is unable to achieve isolated quad set in long seated position. Goal by week 2 is to achieve heel lift with a quad set.  
\*Dosing quad sets: 10 minutes of 10 second squeeze/10 second rest, x5 times a day.
- 4.** Straight leg raises: Ensure quadriceps is activated and is maintaining contraction throughout the SLR range to eliminate extensor lag. Aim for a calf tap and elimination of extensor lag by week 3. Calf tap: the calf taps/skims the table while the heel stays elevated as the leg descends to starting position. Continue doing SLR until 10# ankle weight is achieved.
- 5.** Do not force flexion ROM, but encourage steady progression. Patellar mobility is imperative. Use gentle soft tissue techniques for areas such as anterior interval/fat pad, quadriceps, hamstrings, and scar management. If 90° of flexion is not achieved by week 4, notify physician's office.
- 6.** Start double leg (DL) mini squats and leg press from 0° to 60° initially, then progress to 90° as tolerated. Single leg (SL) activities may be initiated at week 4 with SL leg press and step-ups, then advancing to SL activities as tolerated. Loaded leg extensions are prohibited.  
\*Squat progressions example: DL leg press, DL mini squats, DL chair squats, DL body weight squats, SL leg press, SL step ups, Static lunge split squat, SL step downs, SL squats, SL split squat with elevated back leg, walking lunges, SL sit to stands, SL slide outs.

**Theodore B. Shybut, MD FAANA FAAOS**

Southern California Orthopedic Institute

Knee, Shoulder, Elbow Arthroscopy &

Reconstructive Surgery

Burbank, Valencia / Santa Clarita, Van Nuys

@shybutMD / [www.shybutmd.com](http://www.shybutmd.com)



7. Pre-run/pre-jump program includes tempo-based activities with focus on the deceleration phase such as DL speed squats, DL drop squats, DL “bounce bounce bounce squat”, then progress to alternating SL drop squats. Also, intermittently increase the tempo of regular strengthening exercises to align with the timing requirements of jogging and jumping.

8. Walk/Jog program: MD approval required. Begin on treadmill with 2- 3 days per week. Begin with 1:1 or 2:1 walk to jog ratios, (ie. 1 min walk to 1 min jog or 2 min walk to 1 min jog). Then progress each week by 1 min jog until 12-15 min of jogging is achieved.

9. Plyometric program: MD approval required. Begin with small DL jumps, jump rope, and small depth jump landings& box jumps. Progress box height as skill is mastered. Ensure equal weighted DL take-off and landing before progressing to SL plyometrics. Initiate SL plyometrics with alternating L and R landings in place and then advance to SL hops. Begin a sportsmetric based plyometric program when released by surgeon.

10. Isokinetic protocol: After 16 weeks and with MD approval, may begin training and testing with 300°/sec and progress to 180°/sec. Do not proceed if patient has history of anterior knee pain.

11. Return to Play Progression: a graded re-exposure is essential. Return to non-contact practice, return to contact practice, return to scrimmage, return to interval play, return to full time play.

12. **Prevention of future ACL injury** requires ongoing dedication to correcting functional movement deficits identified during rehabilitation. **FIFA 11+** is an injury prevention program that has demonstrated efficacy. Dr. Shybut recommends athletes, therapists, and trainers utilize this program and incorporate the exercises into their ongoing conditioning.

**Theodore B. Shybut, MD FAANA FAAOS**  
Southern California Orthopedic Institute  
Knee, Shoulder, Elbow Arthroscopy &  
Reconstructive Surgery  
Burbank, Valencia / Santa Clarita, Van Nuys  
@shybutMD / [www.shybutmd.com](http://www.shybutmd.com)



## **PHASE 1 - ACUTE (0-6 Weeks)**

### **PHASE GOALS: PROTECT GRAFT, RESTORE AMBULATION & ADL STATUS RANGE OF MOTION**

0-1 Week – Full extension and progress flexion to 60°

1- 4 Weeks – Maintain full extension and progress to 90°

4+ Weeks – Gradually progress flexion to 120°

### **WEIGHT BEARING**

0-2 Weeks – Toe Touch Weight Bearing

2-4 Weeks – Partial Weight Bearing

4+ Week – FWB with symmetrical gait

### **BRACE & CRUTCH USE**

0-1 Week -- Brace locked in 0° extension

1-2 Weeks -- Unlock brace 0° to 60°

2-4 Weeks – Unlock brace 0° to 90°

4-6 Weeks -- Brace unlocked to full if good quad control and full extension

6+ Weeks -- D/C crutches when gait is normal; D/C brace for home ambulation; continue brace for community ambulation until MD approval to D/C

### **STRENGTHENING**

Quad sets, straight leg raises, hip abduction

2+ Weeks: Pre-gait, TKEs, calf raises

4+ Weeks: Static balance, mini squats, stationary biking: must be > 110° knee flexion

### **CRITERIA FOR FULL AMBULATION**

- > 0 DEG KNEE EXTENSION & 90 DEG KNEE FLEXION
- > 30 STRAIGHT LEG RAISES WITHOUT LAG
- MINIMAL EFFUSION/PAIN, & SYMMETRICAL GAIT WITHOUT A LIMP
- MD OR PT APPROVAL

## **PHASE 2 – STRENGTH (6-12 Weeks)**

### **PHASE GOALS: IMPROVE STRENGTH AND PROTECT THE GRAFT**

[www.sportssurgeonsocal.com](http://www.sportssurgeonsocal.com)

@ShybutMD

661-290-5473

**Theodore B. Shybut, MD FAANA FAAOS**  
Southern California Orthopedic Institute  
Knee, Shoulder, Elbow Arthroscopy &  
Reconstructive Surgery  
Burbank, Valencia / Santa Clarita, Van Nuys  
@shybutMD / [www.shybutmd.com](http://www.shybutmd.com)



### **RANGE OF MOTION**

6+ Weeks – Maintain full extension and progress to full flexion

### **STRENGTHENING**

#### **LOADED FLEXION > 90° IS PROHIBITED**

Leg press, step ups, step downs, RDLs, lunges, Bulgarian squats, wall sits

Squat progression: bodyweight squats → single leg squats

Advance hip abduction & glut strength: band walks, lateral lunge, reverse lunge, bridges, hip thrusters

Core exercises: planks, side planks, v-ups, Russian twist, superman

Balance training: foam pad, balance board, BOSU

### **CONDITIONING**

Initiate dynamic warm-up: frankenstein kicks, leg swings, knee hugs, heel sweeps, heel/toe walks, oil rigs, lateral lunge, hip rotation, inch worm, speed squats

Stationary bike, elliptical, & rowing machine

Swimming: progress kicking gradually and pain-free, no flip turns

### **CRITERIA FOR PHASE 2**

- **PAIN LESS THAN 3/10 (WORST)**
- **WITHIN 2 DEG NORMAL KNEE EXTENSION & 120 DEG KNEE FLEXION**
- **SYMMETRICAL BODY WEIGHT SQUAT**
- **MINIMAL EFFUSION, PAIN, & SYMMETRICAL GAIT WITHOUT A LIMP**

### **PHASE 3 – INITIATE JOGGING AND DOUBLE LEG PLYOMETRICS (12-20 Weeks)**

**PHASE GOALS: IMPROVE STRENGTH AND INITIATE JOGGING PROGRAM**

#### **RANGE OF MOTION**

12+ Weeks – Maintain symmetry & pain-free with overpressure

### **STRENGTHENING**

Leg press, step ups, step downs, RDLs, lunges, Bulgarian squats, wall sits

Squat progression: bodyweight squats → single leg squats

Advance hip abduction & glut strength: band walks, lateral lunge, reverse lunge, bridges, hip thrusters

[www.sportssurgeonsocal.com](http://www.sportssurgeonsocal.com)

@ShybutMD

661-290-5473

**Theodore B. Shybut, MD FAANA FAAOS**

**Southern California Orthopedic Institute**

Knee, Shoulder, Elbow Arthroscopy &

Reconstructive Surgery

Burbank, Valencia / Santa Clarita, Van Nuys

@shybutMD / [www.shybutmd.com](http://www.shybutmd.com)

Core exercises: planks, side planks, v-ups, Russian twist, superman

Balance training: foam pad, balance board, BOSU



**CONDITIONING**

Dynamic warm up & integrate sport specific warm up

Stationary bike, elliptical, swimming, & rowing machine

12+ Weeks -- Treadmill walk/jog progressions: begin with 30"-1' W/J intervals, advance jog time by 1 min each week

16+ Weeks – Double leg jump rope and shuttle jumps

**CRITERIA FOR JOGGING AND DOUBLE LEG JUMP ROPE**

- PAIN LESS THAN 3 / 10 (WORST)
- WITHIN 2 DEG NORMAL KNEE EXTENSION & 120 DEG KNEE FLEXION
- QUADRICEPS & HAMSTRING STRENGTH > 60% NORMAL
- LESS THAN 4cm DEFICIT ON SINGLE-LEG SQUAT (ANTERIOR REACH)
- AT LEAST 1 MINUTE OF SINGLE LEG SQUATS
- MD APPROVAL

**PHASE 4 – STRENGTH, AGILITY, AND PLYOMETRIC (20-24 Weeks)**

**PHASE GOALS: INITIATE DYNAMIC MOVEMENTS & SINGLE LEG PLYOMETRICS**

**RANGE OF MOTION**

20+ Weeks – Promote and maintain symmetry

**STRENGTHENING**

Gym strengthening: squats, deadlifts, initiate olympic lifting

SL strengthening: SL squats, sit to stands, ball slams, step ups/downs

Dynamic core exercises: mountain climbers, planks, pikes, pale off press

Integrate interval strength circuits & work/rest timed intervals

**CONDITIONING**

Dynamic warm up & sports specific warm up

Stationary bike, elliptical, swimming, & rowing machine

Advance to track workouts: jog straights & walk curves

[www.sportssurgeonsocal.com](http://www.sportssurgeonsocal.com)

@ShybutMD

661-290-5473

**Theodore B. Shybut, MD FAANA FAAOS**  
Southern California Orthopedic Institute  
Knee, Shoulder, Elbow Arthroscopy &  
Reconstructive Surgery  
Burbank, Valencia / Santa Clarita, Van Nuys  
@shybutMD / [www.shybutmd.com](http://www.shybutmd.com)



**PLYOMETRICS & AGILITY:**

Ladder drills, footwork agility drills, cone drills

Double leg plyos: jump rope, line jumps, cone jumps, depth jumps, box jumps

Single leg landings: alternating, line jumps, hops

High intensity predictable patterned movements, incorporate sport specific drills

**CRITERIA FOR PLYOMETRICS AND AGILITY  
PAIN LESS THAN 2 / 10 (WORST)**

- **QUAD & HAM STRENGTH > 80% NORMAL; > 50% H/Q RATIO FOR FEMALES**
- **AT LEAST 3 MINUTES OF SINGLE LEG SQUATS (RESISTED)**
- **JOGGING >15 MINUTES ON TREADMILL**
- **< 5 ON LANDING ERROR SCORING SYSTEM (LESS)**
- **MD OR PT APPROVAL**

**PHASE 5 – ADVANCED STRENGTH, AGILITY, AND PLYOMETRIC  
(24-28 Weeks)**

**PHASE GOALS: INTRODUCE POWER MOVEMENTS  
RANGE OF MOTION**

24+ Weeks – Maintain symmetry & pain free overpressure

**STRENGTHENING**

Gym strengthening: squats, deadlifts, olympic lifting

Interval strength circuits & work/rest timed intervals

Dynamic eccentric loading: double & single leg

Dynamic core: rotational and anti-rotational drills

Isokinetic training protocols: begin with 300°/sec, progress to 180°/sec

**CONDITIONING**

Dynamic warm-up

Biking, elliptical, jogging, swimming & rowing

Track workouts: advance to linear speed drills and sprinting drills

**PLYOMETRICS & AGILITY (2-3 DAYS/WEEK):**

Tuck jumps, squat jumps, bounding, SL hop, SL triple hop, SL cross over hop

[www.sportssurgeonsocal.com](http://www.sportssurgeonsocal.com)  
**@ShybutMD**

**661-290-5473**

**Theodore B. Shybut, MD FAANA FAAOS**

**Southern California Orthopedic Institute**

Knee, Shoulder, Elbow Arthroscopy &

Reconstructive Surgery

Burbank, Valencia / Santa Clarita, Van Nuys

@shybutMD / [www.shybutmd.com](http://www.shybutmd.com)

Change of direction drills: begin with < 90°, progress to 90° and greater

Introduce unpredictable agility movements

Non-contact sports specific drills



### **CRITERIA FOR ADVANCED AGILITY & SL PLYOMETRICS**

- **PAIN LESS THAN 2 / 10 (WORST)**
- **QUAD & HAM STRENGTH > 80% NORMAL; > 50% H/Q RATIO FOR FEMALES**
- **AT LEAST 3 MINUTES OF SINGLE LEG SQUATS (RESISTED)**
- **JOGGING >15 MINUTES ON TRACK OR PAVED SURFACE**
- **MD OR PT APPROVAL**

### **PHASE 6 - RETURN TO PLAY (28+ Weeks)**

#### **PHASE GOALS: INITIATE SPORTS SPECIFIC MOVEMENTS & RETURN TO PLAY STRENGTHENING**

Gym strengthening: squats, deadlifts, and olympic lifting

Interval strength circuits & work/rest timed intervals

Dynamic strength and core exercises

Complex movement patterns

Isokinetic protocols: 300°, 180°, and 60°/sec

#### **CONDITIONING**

Jogging, biking, swimming, rowing, & interval sprint workouts

#### **PLYOMETRICS & AGILITY (2-3 DAYS/WEEK)**

Max effort DL and SL jumps → progress with rotation

Lateral & rotational agility drills

Unpredictable cutting agility

Non contract drills → contact drills with MD approval

**Return to practice → return to contact practice → return to scrimmage → return to interval play → return to full play**

#### **CRITERIA FOR RETURN TO PLAY**

- **PAIN LESS THAN 2 / 10 (WORST)**
- **> 75/100 ON ACL-RSI SURVEY**

[www.sportssurgeonsocal.com](http://www.sportssurgeonsocal.com)

@ShybutMD

661-290-5473

**Theodore B. Shybut, MD FAANA FAAOS**  
Southern California Orthopedic Institute  
Knee, Shoulder, Elbow Arthroscopy &  
Reconstructive Surgery  
Burbank, Valencia / Santa Clarita, Van Nuys  
@shybutMD / [www.shybutmd.com](http://www.shybutmd.com)



- **QUAD & HAM STRENGTH > 90% NORMAL; > 60% H/Q RATIO FOR FEMALES**
- **90% NORMAL ON SINGLE-LEG HOP TESTS**
- **95% NORMAL FIGURE OF 8, 5-10-5 PRO-AGILITY, & SL VERTICAL JUMP**
- **MD APPROVAL**